



## IceWind Inter-comparison of Icing Production Loss Models

**Davis, Neil; Hahmann, Andrea N.; Clausen, Niels-Erik; Pinson, Pierre; Zagar, Mark; Byrkjedal, Øyvind ; Karlsson, Timo ; Wallenius, Tomas ; Turkia, Ville ; Söderberg, Stefan**

*Total number of authors:*

11

*Publication date:*

2014

[Link back to DTU Orbit](#)

### *Citation (APA):*

Davis, N. (Author), Hahmann, A. N. (Author), Clausen, N-E. (Author), Pinson, P. (Author), Zagar, M. (Author), Byrkjedal, Ø. (Author), Karlsson, T. (Author), Wallenius, T. (Author), Turkia, V. (Author), Söderberg, S. (Author), & Baltscheffsky, M. (Author). (2014). IceWind Inter-comparison of Icing Production Loss Models. Sound/Visual production (digital)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# IceWind Inter-comparison of Icing Production Loss Models

- DTU Wind Energy
  - Neil Davis
  - Andrea Hahmann
  - Niels-Erik Clausen
  - Pierre Pinson (**DTU Electro**)
  - Mark Žagar (**Vestas**)
- Kjeller Vindteknikk
  - Øyvind Byrkjedal

- VTT
  - Timo Karlsson
  - Tomas Wallenius
  - Ville Turkia
- WeatherTech Scandinavia
  - Stefan Söderberg
  - Magnus Baltscheffsky



# IceWind


- Nordic Project supported by Top-Level Research Initiative (TFI)
- Improved forecast of wind, waves and icing
- 13 Project partners
- Work Package 1: Wind turbine icing

# Observations

- Selected data from 15 wind farms
  - Averaged to wind farm values, not turbine specific
  - 2 years of data (June 2010-June 2012)
  - Observed icing times from automated approach classifying production loss
  - Data removed when turbines not operating optimally



## Power Deviation from Power Curve Using Nacelle Wind Speed




- Note 2 regimes in different wind farms
- Similar results from both years

# WRF model data

Turbine Icing @ 80m

Init: 2012-12-16\_00:00:00  
Valid: 2012-12-17\_17:00:00


- Provided by Vestas at 3 km
- WSM5 microphysics
- 6 hour spin up cycle
- Provided Fields
  - Wind Speed
  - Temperature
  - Pressure
  - 4 Cloud types
  - Precipitation rate
  - Specific humidity
  - Shortwave radiation
  - Longwave radiation





# WRF model data

- 3D Representation of the atmosphere
- Data interpolated to 40, 80, 120, 160 and 200 m AGL




# Production Loss Models




# DTU model


Farm A




Farm A



Farm A




Farm A




- Mixed model
  - Fits separate models for forecast ice / no ice conditions
- Generalized Additive Model
- Utilizes results from WRF and iceBlade
  - IceBlade modified to include cloud ice for WSM5 microphysics
- Fit separately for each farm in this study with consistent variables

# Kjeller Model

- Two-parameter power curve
  - Suggested by wind tunnel results
  - Ice mass and wind speed
  - Tuned and validated using operational data
- Uses a standard cylinder for ice mass modeling
- Assumes power yield of 0 at approximately 9 kg/m




# VTT



- Based on statistical analysis of power loss observations
- Produces an estimate for power loss due to rotor icing
  - Based on wind speed and length of icing event
- Independent of icing or production forecasting methods
- Requires external icing forecast
  - Used iceBlade accumulated icing for this comparison

# WeatherTech Scandinavia

- WICE – WeatherTech Ice Model
- Artificial Neural Network
  - Trained with observed clean & iced production
- Tested for different turbines & locations
- Either forecast or assessment tool




# Results

# Terminology

- 2 Years (defined June – May)
  - Year 1 Used to fit statistical models
  - Year 2 Evaluation year
- 2 Power estimates
  - Gross: power estimate without icing
  - Iced: power estimate with icing
- 2 Observed Conditions
  - Ice: times when observations suggest icing
  - No Ice: times without icing
- 14 Farms (Labeled A-O, ex. G)
- 4 Models (Labeled I-IV)


## Gross Power RMSE



- Power curve fit to nacelle wind speed
- Gross estimates similar across models
- Much larger errors for observed icing cases
- Error pattern similar to impact of icing from boxplot




## Distribution of Gross Power Bias




- Peak near zero for all models
- Symmetrical bias for no ice
- Ice condition skewed positive signifying higher estimated power than observed
- No large deviation across models

## Iced vs Gross Error for Iced Times




- Large improvement in year 1 for Models II, III and IV
- Much smaller improvement in year 2

## Iced vs Gross Bias for Iced Times




- General shift of positive bias to negative
- Year 2 shows larger shift of bias from positive to negative

## Modeled Error for Year 2 Iced Times at Each Wind Farm




- Reasonable performance from all models
- Large differences between models at most sites
- Can pick out sites with low ice impact
- Model III and IV slightly outperform other models at several sites

Model RMSE by Year




- Colors signify years
- Model II appears to have over fit model to year 1
- Models III and IV shows larger errors at sites with less icing than other models

Change in Model Mean Bias (Iced - Gross)



- Using any ice model almost always reduces bias
- Bias reduced more for sites with large amounts of ice
- Not a large change from year to year

Change in Model RMSE (Iced - Gross)



- Iced RMSE often worse than Gross, due to the decrease in performance for non-iced times
- Depending on agreement, bias correction may offset increased error



# Conclusions

- Models perform similarly
- Differences appear mostly due to park conditions
- Large improvements still possible
  - Longer periods for model fit to reduce over fitting
  - WRF runs customized for icing
  - Ice ablation methods & relationship to power
- Agreed upon metric is needed to help improve the models
  - Bias was improved at most sites
  - RMSE was not improved as much
- Using human input could improve these models, need judgment on when to apply them